

FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10

EL 239201145 US
Express Mail Label Number4/28/99
Date of Deposit

Form PTO-1390-MOD (REV 10-96)		U S Department of Commerce Patent and Trademark Office	ATTORNEY'S DOCKET NUMBER 4-21101/A / PCT
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 09/284615
INTERNATIONAL APPLICATION NO. PCT/EP97/05897	INTERNATIONAL FILING DATE 24 October 1997 (24.10.97)	PRIORITY DATE CLAIMED 28 October 1996 (28.10.96)	
TITLE OF INVENTION METHOD FOR THE OLIGOMERISATION OF PEPTIDES			
APPLICANT(S) FOR DO/EO/US ALEXEY ET AL.			

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau. (See Form PCT/IB/308)
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☐ An executed Declaration and Power of Attorney (original or copy) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included.

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Sequence listing and sequence diskette; Unsigned declaration.

17. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492(a) (1)-(5)):

Search Report has been prepared by the EPO or JPO \$840

International preliminary examination fee paid to USPTO (37 CFR 1.482) \$670

No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)). \$760

Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO. \$970

International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4). \$96

ENTER APPROPRIATE BASIC FEE AMOUNT =

Surcharge of **\$130** for furnishing the oath of declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	33	- 20 =	13	X \$	18
Independent claims	3	- 3 =	0	X \$	78
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				+	\$ 260
TOTAL OF ABOVE CALCULATIONS =					\$ 1,074
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).					\$
SUBTOTAL =					\$ 1,074
Processing fee of \$130 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).					+
TOTAL NATIONAL FEE =					\$ 1,074
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40 per property					+
TOTAL FEES ENCLOSED =					\$ 1,074
					Amount to be: refunded
					charged

CALCULATIONS PTO USE ONLY

- a. ☐ A check in the amount of \$_____ to cover the above fees is enclosed.
- b. ☒ Please charge Deposit Account No. 19-0134 in the name of Novartis Corporation in the amount of \$1,074 to cover the above fees. A duplicate copy of this form is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 19-0134 in the name of Novartis Corporation.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

Send all correspondence to the address associated with Customer No. 001095, which is currently:

Michael W. Glynn
Novartis Corporation
Patent and Trademark Dept.
564 Morris Avenue
Summit, NJ 07901-1027

Myra H. McCormack, Ph.D.
Attorney for Applicants
Reg. No. 36,602
(908) 522-6938

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN RE PCT NATIONAL STAGE APPLICATION OF

ALEXEY ET AL.

INTERNATIONAL APPLICATION NO: PCT/EP97/05897

FILED: 24 OCTOBER 1997

U.S. APPLICATION NO: Not Yet Known

35 USC §371 DATE: Herewith

FOR: METHOD FOR THE OLIGOMERISATION OF PEPTIDES

Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Prior to the examination of the above-referenced application, kindly amend the application as follows:

IN THE SPECIFICATION

Kindly replace the existing Sequence Listing with the Substitute Sequence Listing provided herewith as pages 19-25.

IN THE CLAIMS

Kindly renumber the claims pages beginning with page 26.

At the top of the first page of claims kindly delete "Claims" and insert therefor --What is Claimed is--.

Kindly amend the following claims:

1. (Amended) An oligomer comprising at least 2 [or more than 2] units wherein each unit comprises a peptidic domain capable of oligomerizing and a domain capable of binding to an acceptor, wherein the oligomerizing domain is not an antibody or a functional antibody fragment from [the] a constant region.
6. (Amended) An oligomer according to claim 1, wherein the peptidic domain capable of oligomerizing and the domain capable of binding to an acceptor are connected via a spacer[(hinge region)].

14. (Amended) The oligomer of [Use of an oligomer according to] claim 1 wherein the oligomer is useful for the identification and/or marking of acceptors.

15. (Amended) The oligomer of [Use of an oligomer according to] claim 1 wherein the oligomer can bind to [for] eukaryotic cells, bacteria or viruses [targeting].

16. (Amended) The oligomer of [Use of an oligomer according to] claim 1 wherein the oligomer can bind to a cell [for cell targeting].

17. (Amended) The oligomer of [Use according to] claim 16 wherein the oligomer can bind to [for targeting of] B-cell lymphomas.

18. (Amended) The oligomer of [Use of an oligomer according to] claim 1 wherein the oligomer can inhibit [for the inhibition of] protein-protein interactions.

19. (Amended) The oligomer of [Use of an oligomer according to] claim 12 wherein the oligomer is [as] a chelating agent.

20. (Amended) The oligomer of [Use of an oligomer according to] claim 12 wherein the oligomer is [as] a crosslinking agent.

21. (Amended) The oligomer of [Use of oligomers according to] claim 1 wherein the oligomer is used in [for] the construction of libraries.

22. (Amended) The oligomer of [Use of oligomers according to] claim 1 wherein the oligomer induces [for the induction of] apoptosis.

23. (Amended) The oligomer of [Use of oligomers according to] claim 1 wherein the oligomer is capable of inhibiting [for the] intracellular [inhibition of] transcription factor binding, gene regulating molecules and/or enzymatic activities.

24. (Amended) The oligomer of [Use of oligomers according to] claim 1 wherein the oligomer inhibits [for prevention of] tumor metastatization.

25. (Amended) The oligomer of {Use of oligomers according to} claim 1 wherein the oligomer is used [in vitro as one of the binding reagents] in an enzyme immunoassay.

26. (Amended) The oligomer of [Use of oligomers according to] claim 1 wherein the oligomer is used [in vitro as one of the binding reagents] in a radioimmunoassay[s].

31. (Amended) A method for the production of an oligomer [according to claim 1] comprising the steps of:

oligomerizing a unit, wherein each unit comprises at least one peptidic domain capable of oligomerizing and at least one domain capable of binding to an acceptor, wherein the peptidic domain is not an antibody or a functional antibody fragment from a constant region of an antibody; and,

isolating the oligomer produced therefrom.

Kindly add the following claim:

33. (New). The oligomer of claim 6 wherein the spacer comprises at least one hinge region.


REMARKS

Claims 1, 6, 14-26 and 31 have been amended. Claim 33 has been add and incorporates material from claim 6 as originally filed.

A Substitute Sequence Listing has been provided along with a Computer Readable Form of the Sequence Listing. The under signed hereby states that the Paper Copy and the Computer Readable Form, submitted in accordance with 37 CFR 1.821 including 1.821(c) and 1.821(e) are identical. No new matter has been added by any of these amendments. Favorable consideration of this application is respectfully requested.

Respectfully submitted,

Novartis Corporation
Patent and Trademark Dept.
564 Morris Avenue
Summit, NJ 07901-1027
(908) 522-6938


Myra H. McCormack, Ph.D.
Attorney for Applicants
Reg. No. 36,602

Date: April 28, 1999

SEQUENCE LISTING

□

□

<110> Terskikh, Alexey

□

Crameri, Reto

□

Mach, Jean-Pierre

□

Le Doussal, Jean-Marc

□

Kajava, Andrey

□

□

<120> Method for the Oligomerisation of Peptides

□

□

<130> 4-21101/A

□

□

<140> unknown

□

<141> 1999-04-28

□

□

<150> PCT/EP 97/05897

□

<151> 1997-10-24

□

□

<160> 13

□

□

<170> PatentIn Ver. 2.0

□

□

<210> 1

□

<211> 12

□

<212> PRT

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:peptide

□

□

<400> 1

□

Ser Val Trp Arg Trp Leu Pro Tyr Asp Lys Tyr Glu

□

1

5

10

□

□

□

<210> 2

□

<211> 10

□

<212> PRT

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:peptide

□

□

<400> 2

□

Ala Asp Gly Ala Cys Arg Asn Pro Trp Cys

□

1

5

10

□

□

<210> 3

□

<211> 19

□

<212> PRT

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:peptide

□

□

<400> 3

□

Thr Ala Ala Gly Leu Cys Glu Phe Asp Gln Ala Leu Leu Arg Tyr Thr

□

1

5

10

15

□

□

Cys Pro Thr

□

□

□

□

<210> 4

□

<211> 41

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 4

□

agatcctcga ggggtggagac tgctgcccac agatgcttag a

41

□

□

<210> 5

□

<211> 26

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 5

□

gcactagtag aaccgccacc cggggt

26

□

□

<210> 6

□

<211> 39

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 6

□

gatccgctga cggcgcttgc cgtaccccgt ggtgcggtc

39

□

□

<210> 7

□

<211> 39

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 7

□

tcgagaccgc accacgggtt acggcaagcg ccgtcagcg

39

□

□

<210> 8

□

<211> 36

□

<212> DNA

□

<213> Artificial Sequence

□

□
 <220>
 □
 <223> Description of Artificial Sequence:PCR primer
 □

□
 <400> 8
 □
 gatccactgc tgcaggtctg tgcgaatccg accagc 36
 □

□
 <210> 9
 □
 <211> 36
 □
 <212> DNA
 □
 <213> Artificial Sequence
 □

□
 <220>
 □
 <223> Description of Artificial Sequence:PCR primer
 □

□
 <400> 9
 □
 tcgagctggg cgaattcgca cagacctgca gcagtg 36
 □

□
 <210> 10
 □
 <211> 42
 □
 <212> DNA
 □
 <213> Artificial Sequence
 □

□
 <220>
 □
 <223> Description of Artificial Sequence:PCR primer
 □

□
 <400> 10
 □
 gatacctctgt ttggcggttg ctgccgtacg acaaatacga ac 42
 □

□

<210> 11

□

<211> 42

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 11

□

tcgagttcgt atttgtcgta cggcagccaa cgccaaacag ag

42

□

□

<210> 12

□

<211> 78

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 12

□

tcgagcggca gccgcagccg aaaccgcagc cgcagccgca gccgcagccg aaaccgcagc 60

□

cgaaaccgga accggaag

78

□

□

<210> 13

□

<211> 78

□

<212> DNA

□

<213> Artificial Sequence

□

□

<220>

□

<223> Description of Artificial Sequence:PCR primer

□

□

<400> 13

□

tcgacttccg gttccggttt cggctgcggt ttcggtgcg gctgcggctg cggctgcggt 60

□

ttcggtgcg gctgcggc

78

□

□

□

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151

METHOD FOR THE OLIGOMERISATION OF PEPTIDES

Great strides made over the past years in understanding of molecular interactions in the realm of biomolecules such as proteins and nucleic acids benefited from the isolation of artificial polypeptide "ligands" with *de novo* binding activities to "receptors". A powerful means of evolving artificial ligands is offered by the use of large polypeptide libraries, displayed on the surface of filamentous bacteriophage as a fusion with the phage coat proteins. In particular, isolation of new peptide ligands allowed, for example, to map antibodies binding sites, to find an important residues in HLA-DR molecules, to identify proteases substrates and inhibitors. Obviously, isolation of new peptide ligands with high affinity towards their target "receptors" can help to answer important biological questions and is of great interest for pharmaceutical development. However, apart from some exceptions, only low affinity (micromolar range) ligands were isolated from peptide libraries. This can be readily explained by a high degree of conformational freedom and a few number of contact residues within a short peptide molecule.

In order to improve this situation, a new type of high avidity binding molecule is constructed by harnessing the effect of multivalent interaction. In the current examples a short peptide ligand is expressed *via* a semi-rigid hinge region as a fusion with a coiled coil domain from Cartilage Oligomeric Matrix Protein (COMP), resulting in a pentameric multivalent binding molecule. In the case of a Pentabody (Pab-S) as described here, a peptide ligand (S) specific for the mouse B-cell lymphoma (BCL₁) surface idiotype, is selected from a phage displayed library. Pab-S molecule specifically bind the BCL₁ cells surface idiotype with an apparent avidity of about 1nM, which corresponds to a 2×10^5 fold increase, compared to the affinity of the synthetic peptide S itself. Equilibrium binding studies strongly suggest that high avidity of Pab-S is a result of multivalent interaction of its "peptide heads" and surface BCL₁ immunoglobulin receptor. As demonstrated by gel filtration chromatography, SDS-PAGE analysis and circular dichroism, Pab-S is a stable homopentamer of about 85kDa, with the five chains crosslinked by disulfide bridges. Pab-S can be reversibly denatured, reconstituting upon renaturation its pentameric structure and full binding activity, providing an easy way to bring different peptide specificity within the same heteropentamer, thus bispecific or multispecific pentabodies.

The inventive oligomers have various unique features.

- Target molecule specificity of inventive oligomer can be provided by a short peptide ligand, representing a "minimal" binding domain, where the primary structural information (i.e. the amino acid sequence) is sufficient for recognition. This demonstrates for the first time, that a low affinity peptide ligand (though specific) can be used to create a high avidity binding molecule.
- A hinge region which dictates the geometry and the dynamic features of multivalent interaction can be easily introduced.
- The inventive oligomers can be produced without extensive post translational modifications and, hence, can be produced easily in microorganisms like *E. coli*.
- The peptide ligands can be attached to the C- or N-terminus of the oligomerizing domain, thus providing various binding geometry, with maximal distance between two peptide ligands (for example some attached to the C-terminus and some to the N-terminus in one oligomer).
- If coiled coil α -helical domains are used, the resulting oligomers are in most cases well soluble due to the intrinsic structure.
- The COMP domain can be used generally for oligomerization of compounds bound thereto.
- It is a general method for the oligomerization of peptides.

Detailed description of the invention

The present invention concerns an oligomer comprising 2 or more than 2 units, wherein each unit comprises a peptidic domain capable of oligomerizing and a domain capable of binding to an acceptor (ligand), wherein the oligomerizing domain is not an antibody or a functional antibody fragment from the constant region.

A functional antibody fragment is, for example, the F_c domain or the constant region of the light or heavy chain.

In a preferred embodiment of the invention the inventive oligomer comprises more than 2 preferably more than 4 units and most preferably the inventive oligomer consist of 5 units.

In another preferred embodiment the individual units oligomerize spontaneously. A peptidic domain capable of oligomerizing is, e.g. a peptide that is known for its tendency to

oligomerize like a pentamerization domain of the Cartilage Oligomeric Matrix Protein (Slavin & Strober (1978) *Nature* **272**, 624-626). Especially preferred units can be separated from the oligomer and re-oligomerized without loss of activity.

In a further preferred embodiment of the invention each of monomeric units has less than 600 amino acids.

The acceptor, to which the inventive oligomer binds, may be of various origin like, for example, any protein that specifically binds to a partner like a substrate, an inhibitor, an activator, an antibody, a receptor and the like. A preferred acceptor is, for example, an antibody or a receptor. Examples for such preferred antibodies or receptors are BCL₁ cells from a B-cell lymphoma surface idiotype or any other receptor expressed on cell surface.

The inventive oligomer may comprise units having the same or different specificities in order to bind to the same or to distinct acceptors; wherein the domain capable of binding to an acceptor can be attached to the C- and/or N-terminus of the oligomerizing domain.

Embraced by the scope of the present invention is also the individual unit capable of oligomerizing as described above, a method for the synthesis of this unit, the expression vector used for this synthesis, as well as the host comprising said expression vector. Preferred hosts are of microbiological origin like *E. coli*.

There are several ways to synthesize the inventive units.

- Via genetic engineering: An expression vector is constructed comprising an expression cassette for the complete inventive unit, that is expressed in a suitable host. Suitable expression cassettes can be prepared by standard techniques in genetic engineering. Beside the information necessary for the synthesis of the desired unit, the expression cassette may additionally contain a signal sequence that forces the secretion of the protein produced. It is also possible to express the inventive unit as a fusion protein with phage coat protein, especially those from filamentous phages and phagemids
- Chemically: Due to the relatively short length of the individual units, they can be synthesized chemically, e.g. via solid phase synthesis.
- Mixed: A part of the inventive unit is expressed by a suitable host and is connected with a chemically synthesized part. For example, the oligomerizing part is synthesized by a

microorganism and is elongated by chemical synthesis to add a domain capable of binding to an acceptor.

The peptidic domain capable of oligomerizing and the domain capable of binding to an acceptor may be connected directly or via a spacer (hinge region) to provide, for example, a greater flexibility of the binding domain. For example, a proline-rich sequence of a spacer is supposed to prevent formation of secondary structure elements and as a consequence a fixed 3-D structure. Moreover, a spacer region like that is generally supposed to be quite rigid due to the conformational constraints of the proline residues, bringing a beneficial effect for the cooperativity of the multivalent binding. Accordingly, in another preferred embodiment of the invention, the peptidic domain capable of oligomerizing and the domain capable of binding to an acceptor are connected via a spacer, that comprises preferably a proline-rich region.

Furthermore, the C-terminus of some or all units of the oligomer may be modified by the addition of a further functional domain, like a marker, an enzymatic or a cytotoxic domain, or a domain that adds additional binding properties, for example to metal atoms or other structural compounds, that can be used, e.g., in affinity chromatography.

The individual units may be connected via disulfide bonds, e.g. spontaneously, or via one or more linker molecules. Such linker molecules are molecules bearing two or more reactive groups like -SH, -N₃, -COOH, -COBr, -COCl, -NH₂, -CHO, -CO-O-CO-, -CO-NH-CO-. Examples are N-5-azido-2-nitrobenzoyloxysuccinimide, p-azidophenazylbromide, p-azidophenyl glyoxal, N-4-(azidophenylthio)phthalimide, bis(sulfosuccinimidyl) suberate, bis-maleimido-hexane, bis[2-(succinimidooxycarbonyloxy)ethyl] sulfone, 1,5-difluoro-2,4-dinitrobenzene, 4,4'-diisothiocyano-2,2'-disulfonic acid stilbene, dimethyl adipimidate, dimethyl pimelimidate, dimethyl suberimidate, dithiobis(succinimidylpropionate), disuccinimidyl suberate, disuccinimidyl tatrane, dimethyl 3,3'-dithiobispropionimidate, 4,4'-dithiobisphenylazide, 3,3'-dithiobis (succinimidylpropionate), ethyl-4-azidophenyl-1,4-dithiobutyrimidate, 1-azido-4-fluoro-3-nitrobenzene, N-hydroxysuccinimidyl-4-azidobenzoate, methyl-4-azidobenzoimidate, m-maleimidobenzoyl-N-hydroxysulfo-succinimide ester, N-hydroxysuccinimidyl-4-azidosalicylic acid, p-nitrophenyl-2-diazo-3,3,3-trifluoro propionate, N-succinimidyl(4-azidophenyl)-1,3'-dithiopropionate, sulfosuccinimidyl 2-(m-azido-o-nitrobenzamido)-ethyl-1,3'-dithiopropionate, N-succinimidyl-6(4'-azido-2'-nitrophenyl-amino)hexanoate, sulfosuccin-

imidyl 2-(p-azidosalicylamido)ethyl-1,3'-dithiopropionate, N-succinimidyl(4-iodoacetyl)aminobenzoate, succinimidyl 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate, succinimidyl 4-(p-maleimidophenyl)-butyrate, N-succinimidyl 3-(2-pyridyldithio)propionate, bis[2-(sulfosuccinimidooxy-carbonyloxy)ethyl]sulfone, disulfosuccinimidyl tatrane, ethylene glycol-bis(sulfosuccinimidylsuccinate), m-maleimidobenzoyl-N-hydroxysulfosuccinate, sulfosuccinimidyl (4-azidophenyldithio)-propionate, sulfosuccinimidyl 6-(4'-azido-2'-nitrophenylamino)hexanoate, sulfosuccinimidyl (4-iodoacetyl)aminobenzoate, sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate, sulfosuccinimidyl 4-(p-maleimidophenyl)butyrate and 2-iminothiolane.

Preferably, oligomerizing units capable of self-assembling are used. The linker molecules mentioned above may be also used additionally to stabilize the oligomer.

As shown in the example, spontaneous pentamerization of COMP assembly domain does not depend on disulfide bond formation, therefore, a preferred embodiment of the inventive oligomer represents a self-assembling molecule able to oligomerize *in vivo* and/or *in vitro*. Thus oligomerization of short peptides bypass folding problems and overcomes expression difficulties previously experienced during oligomerization of relatively complex proteins such as single chain Fv fragments. Moreover, a self-assembling of an oligomer allows it to undergo a reversible denaturation. Thus, heterooligomers can be readily obtained by mixing the inventive units with different specificity (i.e. with different peptide ligands) under denaturing conditions followed, e.g., by dialysis against any physiological buffer. This intrinsic property of the inventive oligomers opens an easy way to produce a chelating oligomer where two or more peptides directed against different epitopes on the same target molecule would be assembled together into a heteropentameric oligomer, allowing to benefit from the power of chelating effect, as it is recently demonstrated for the single chain Fv fragments. The same procedure can be used to produce heterooligomers with affinity to two or more receptors.

Binding studies of ¹²⁵I labeled Pab-S on BCL1 cells, for example, reveal an avidity of about 1 nM for the surface idiotype which is 2×10^5 fold higher compared to IC₅₀ of the peptide S itself (200 μM). Pab-S can be reversibly denatured (4M, urea 95°C, 15 min), regaining full binding activity after renaturation by dialysis in PBS. Data show that Pab-S has remarkable thermostability, retaining the binding activity even after autoclavation (120°C, 20 min).

Furthermore, it is expected that an oligomer harboring a peptide ligand with a nanomolar intrinsic affinity may attain a femtomolar range of avidity, approaching that of streptavidin-biotin interaction

Based on the properties of the inventive compounds, there are several applications for the inventive oligomers, that are all part of the invention.

- Based on the equilibrium binding results of the example Pab-S it is evident that inventive oligomers are excellent molecules for eukaryotic cell, bacteria or viruses targeting; and especially for cell targeting. As it is shown for Pab-S, the avidity for the cell-surface immobilized receptor is much greater than the avidity for the soluble form of the same receptor. This property allows, for example, efficient targeting of the B-cell lymphomas, even in the presence of relatively high level of circulating idiotype antibody.
- Based on the major feature of high avidity binding, especially for the surface immobilized antigens, the inventive oligomers can also be used as an inhibitor of protein-protein interactions, especially those occurring on the cell surface and in solution.
- Heteropentamers can be produced by reassociating two or more different units of the inventive oligomers together. These heterodimers can bind to different epitopes on the same or different molecules and, e.g., can be used as a chelating and/or crosslinking agent.
- The C-terminus of the molecules can be modified easily, e.g. by introducing peptide ligands with second specificity, a cytotoxic tail, a marker to identify acceptors, or a His-tail to chelate different toxic drugs (e.g. heavy metals) in order to deliver them to the target cells where they can be internalized, via an internalized surface receptor (e.g. immunoglobulin receptor) and kill the target cells.
- It is possible, for example, to isolated a panel of C1q specific peptide ligands which can be used to produce a bispecific form of the inventive oligomers capable of C1q fixation and complement activation or complement inhibition.
- Fusion of the peptide ligands to an Fc receptor results in a Nt-Ct bispecific form of the inventive oligomers capable of imitating the function of an Fc region from immunoglobulin.
- Frameworks of the inventive oligomers can also be used to create random peptide libraries displayed on the filamentous bacteriophages. This allows rapid selection of the

phage displayed inventive oligomers which then can be produced in a soluble form, in a suitable host.

- Due to the relatively short length of the monomer polypeptide chain, the inventive oligomers can be synthesized chemically (e.g. standard Fmoc peptide chemistry, synthesis starts from the C-terminus). The N-terminal position of the peptide ligands allows to synthesized first the core molecule (e.g. the pentamerization domain and a linker) and then split the sample and continue synthesis with different peptide sequences on the N-terminus.
- The same principle of chemical synthesis can be used to produce synthetic peptide library displayed on the inventive oligomers. Namely, synthesizing the N-terminal peptides at random (e.g. by Fmoc chemistry as described above), one can obtain a library of the peptides displayed on the inventive oligomer framework. This library can be absorbed on the target molecules or cells, eluted, purified (e.g. using 6xHis tag) and the N-terminal peptide can be sequenced.
- Furthermore, a framework of the inventive oligomers can be used to oligomerize peptides representing known B-cell epitope in order to obtain an efficient vaccines. Addition of different peptides known to enhance the immune response (such as T-helper epitopes or CR binding peptides derived from C3d) is also possible.
- The inventive units can be expressed directly in a genetically engineered multicellular organism, in order to provide an *in vivo* production of the inventive oligomers.
- The inventive units can be expressed *in vitro* using established transcription-translation systems.
- The inventive units may be used *in vitro* as one of the binding reagents in enzyme immunoassays.
- The inventive units can be used to induce apoptosis.
- Due to the solubility and the high specificity of the compounds they can be used in gene therapy, e.g., to target certain receptors for activation or deactivation; or, if connected to toxic compounds, for their destruction.
- Genes encoding the inventive units may be intracellularly expressed in order to inhibit the binding of transcription factors or gene regulating molecules as well as inhibit enzymatic activities.
- The inventive units may be used to inhibit enzymatic activity or adhesion in order to prevent tumor metastization.

- Furthermore, the epitopes of the inventive units can be used for vaccine development.
- Due their high tendency for oligomerization the Cartilage Oligomeric Matrix Protein can be used generally for the pentamerization of low molecular weight compounds or peptides that are not part of the Cartilage Oligomeric Matrix Protein.

Examples

Peptide synthesis. Peptides are synthesized using standard Fmoc solid phase chemistry on an Applied Biosystems 431A peptide synthesizer, lyophilized, re-dissolved in 50% acetic acid, purified by gel filtration through a G-25 sephadex column and analyzed by mass spectroscopy. For competition studies peptides are dissolved in PBS, adjusted to neutral pH, and peptide concentrations are determined by the method of Waddell based on an absorption difference at 215 and 225 nm.

Bacterial strains. *E. coli* TG1 (Wertman *et al.*, (1986) *Gene* 49, 253-262) is used for propagation of plasmids and phages and *E. coli* SG13009 (QIAEX) is used for production of fusion proteins.

Labeling. Typically, Pab-S (17 µg, 0.2 nmol), B1 IgG (75 µg, 0.5 nmol) or B1 Fab' (50 µg, 1 nmol) are labeled in PBS with 100 µCi (1 µCi = 37 MBq) of ¹²⁵I in a Iodo-Gen (Biorad, 10 µg) coated tubes for 20 min (2 h for Pab-S) at 4°C. Uncoupled iodine is removed by gel filtration on a PD-10 column (Pharmacia). About 40% of the radioactivity for Pab-S and 70% for B1 IgG and B1 Fab' is recovered. Specific activity ranges from 70 to 200 µCi/nmol.

RIA competition on plastic. Polyvinil 96-well plates are coated with BCL1 IgM at 3 µg / ml in PBS for 16 h at 4°C and blocked with 2% MPBS for 1 h at 37°C. Different amounts of peptides are incubated with 10 nCi of ¹²⁵I labeled B1 IgG for 1 h at 37°C. The wells are washed with PBS containing 0.1% of Tween-20 and the amounts of radioactivity retained is measured with an automated multichannel Cobra II γ-counter.

Example 1: Selection of peptides specific for BCL1 surface immunoglobulin receptor.

Cells and antibodies. The BALB/c derived B cell lymphoma BCL₁ (Efimov *et al.*, (1994) *FEBS Letters* 341, 54-58), and the mouse hybridoma B1, secreting an anti-idiotypic monoclonal IgG₁ antibody (B1 IgG), are kindly provided by Dr. Kris Thielemans (Medical

School, VUB, Brussels, Belgium). BCL₁ cells are propagated in BALB/c mice using i.p. injection of 10⁶ cells. The BCL₁ soluble IgM is purified from the serum of a splenomegaly mouse by precipitation with 50% ammonium sulfate, followed by anion exchange and gel filtration chromatography (Mono Q and Superdex 200, Pharmacia). The B1 IgG is purified by protein G-Sepharose (Pharmacia). Fab' fragments are obtained by limited digestion with pepsin, followed by reduction and alkylation, using standard methods.

Peptide selection. Peptide ligands specific for mouse B-cell lymphoma (BCL₁) idiotype, selected from three different peptide libraries displayed on two filamentous bacteriophages libraries of about 10⁷ independent members displaying random hexapeptides called Smith (Scott & Smith (1990) *Science*, **249**, 386-390), and Doorbar (Doorbar & Winter (1994) *J. Mol. Biol.*, **244**, 361-369), are used. In addition a combinatorial library of about 10¹² independent members displaying a tandem of random decapeptides, called Fisch (Fisch *et al.*, *Proc. Natl. Acad. Sci. USA* (1996), **93**, 7761-7766), is used.

Screening of the phage display libraries is performed essentially as described by Fisch *et al.*, (*Proc. Natl. Acad. Sci. USA* (1996), **93**, 7761-7766). Briefly, the purified BCL₁ IgM is coated on immunotubes (Nunc) at 50 µg/ml in PBS (50 mM phosphate, 150m M NaCl, pH 7.4). About 10¹² colony forming units (c.f.u.) of phage from each library are used for each round of selection. After three rounds of panning, individual clones are isolated, and binding of phage to BCL₁ IgM coated on plastic (3µg/ml) is measured by ELISA (Barrett *et al.*, (1992) *Anal. Biochem.* **204**, 357-364) using horseradish peroxidase conjugated anti-M13 antibody (Pharmacia). Plates coated with mouse IgG or with human IgM, are used as negative controls. Specific inhibition of phage binding to BCL₁ IgM, is performed by addition of B1 IgG at 100 µg/ml. The DNA fragments encoding the selected peptides are amplified by PCR and sequenced as described Fisch *et al.*, (*Proc. Natl. Acad. Sci. USA* (1996), **93**, 7761-7766).

Specific phages are selected on the purified BCL₁ IgM, individual phage clones are isolated, and the DNA fragments encoding peptides are amplified by PCR and sequenced (Table 1). Interestingly, two distal Cys residues, potentially capable of forming a disulfide bonded loop, are found in the hexapeptide selected from the Smith library. Idiotype specificity of selected

peptides is demonstrated by inhibition of phage binding with the anti-idiotypic antibody, B1 IgG.

Table 1 Amino acid sequences of selected idiotypic specific peptide ligands for BCL1 immunoglobulin receptor.

Library / sequence [*]	Selected peptides [†]	Sequ ID	name
Doorbar / X6 - PNDKYE - g3p	SVWRWLPYDKYE [‡]	1	D
Smith / ADGA - X6 - g3p	ADGACRNPWC	2	S
Fisch / X10 - ALLRY - X10 - g3p	TAAGLCEFDQALLRYTCPT [¶]	3	F

^{*} Names and amino acid sequences of the phage displayed peptide libraries; X6 and X10 denotes random hexa- and deca-peptide correspondingly.

[†] Peptides selected from random libraries are in bold type, framework residues are in normal type, corresponding synthetic peptides are in italic type.

[‡] An unusual N to Y substitution in the constant part of the peptide from Doorbar library is observed.

[¶] First ten amino acid of the peptide selected from Fisch library was synthesized.


Synthetic peptides corresponding to the selected sequences are prepared as outlined in Table 1. Cysteins containing peptide S is completely oxidized on air losing two hydrogen atoms, as shown by Mass Spectroscopy, which is consistent with the cyclization of the peptide. All synthetic peptides are tested for binding inhibition of ¹²⁵I labeled B1 IgG to BCL₁ IgM coated on plastic. The D and the S peptides inhibit binding with an IC₅₀ of about 60 μM and 200 μM, respectively. No inhibition is seen with peptide F up to 2mM, which is used, thereafter, as a negative control. Interestingly, substitution of Cys residues by Ser in peptide S results in a 10 fold loss of affinity, suggesting that disulfide bounded turn conformation of the peptide is favorable for binding to BCL₁ IgM idiotypic.

Example 2: Pentabody. Molecular design and gene construct

Pentabody fusion protein, is designed consisting of four distinct parts.

- First, a selected peptide specific for BCL₁ IgM idiotypic represented the N-terminal binding domain.

- Second, a 24 amino acid sequence selected from a long camel Ig hinge region (Hamers-Casterman *et al.*, (1993) *Nature* **363**, 446-448) is placed to provide a space necessary for multivalent binding.
- Third, the hinge region is followed by a 55 amino acid long pentamerization domain, a modification of the described coil-coiled COMP assembly domain (Slavin & Strober (1978) *Nature* **272**, 624-626), known to spontaneously form a five-stranded α -helical bundle. Two cystein residues present at the C-terminal of the wild type sequence allowed the formation of inter-chain disulfide bridges. Based on the recently described model of COMP assembly domain (Kajava (1996) *Proteins* **24**, 218-226) two substitutions Lys29Cys and Ala30Cys are introduced in the modified version, making a possible the formation of additional disulfide bonds near the N-terminal part of the assembly domain in order to further stabilize the α -helical bundle.
- Forth, a sequence encoding six histidine residues is placed at the C-terminus of the fusion molecule to facilitate the protein purification *via* the metal-chelating affinity chromatography.



The chimeric genes coding for the different Pab are constructed as designed (see below) and named Pab-S, Pab-D and Pab-F, where the last letters stay for the S, D, and F peptide, respectively (see table 1).

Essentially, the DNA sequences encoding the BCL₁ idiotype specific peptides, as well as the hinge region, are assembled from oligonucleotide duplexes. The COMP pentamerization domain is amplified by PCR simultaneously introducing point mutations, and the fusion genes are assembled into a modified pDS-78 expression vector.

Plasmid construction. The construction of plasmid p3bCOMP encoding a 64 amino acid COMP assembly domain is described in (Slavin & Strober (1978) *Nature* **272**, 624-626). All further constructs are made using standard methods of DNA manipulation (Sambrook *et al.*, (1990) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY)). The DNA fragment encoding the COMP domain (residues 26-80, amino acids numbering as in Slavin & Strober (1978) *Nature* **272**, 624-626) is amplified by PCR from

p3bCOMP template using COMP-Xho-BACK (5'-AG ATC CTC GAG GGT GGA GAC TGC TGC CCA CAG ATG CTT AGA; SEQ ID NO 4) and COMP-Spe-FOR (5'GC ACT AGT AGA ACC GCC ACC CGG GGT; SEQ ID NO 5) primers. Thus the resulting product contains two amino acid substitutions Lys29Cys and Ala30Cys as well as XhoI and SpeI sites at the 5' and 3' end, respectively. The DNA duplexes encoding the peptides S, F, and D, are prepared by annealing the oligonucleotides S-up (5'GA TCC GCT GAC GGC GCT TGC CGT ACC CCG TGG TGC GGT C; SEQ ID NO 6) with S-low (5'TC GAG ACC GCA CCA CGG GTT ACG GCA AGC GCC GTC AGC G; SEQ ID NO 7), F-up (5'GA TCC ACT GCT GCA GGT CTG TGC GAA TCC GAC CAG C; SEQ ID NO 8) with F-low (5'TC GAG CTG GTC GAA TTC GCA CAG ACC TGC AGC AGT G; SEQ ID NO 9) and D-up (5'GA TCC TCT GTT TGG CGT TGG CTG CCG TAC GAC AAA TAC GAA C; SEQ ID NO 10) with D-low (5'TC GAG TTC GTA TTT GTC GTA CGG CAG CCA ACG CCA AAC AGA G; SEQ ID NO 11). All three duplexes contain BamHI and XhoI cohesive ends on 5' and 3' end, respectively. By means of three-part ligation, the duplexes encoding peptides and the PCR amplified COMP domain, restricted with XhoI and SpeI enzymes, are joint together in the pDS78 vector, linearized with BamHI and SpeI enzymes, in front of a 6 histidine tail present in the vector (Stuber *et al.*, in *Immunological Methods*, eds. Lefkovits, I. & Perris, B. (Academic Press, New York), pp.121-152). This generated pSC6H pDC6H and pFC6H plasmids which encode S, D and F peptides respectively. A DNA fragment encoding the 24 amino acid hinge region $[(PQ)_2PK(PQ)_4PKPQPK(PE)_2]$, called PX, is prepared by annealing of PX-up (5'TC GAG CGG CAG CCG CAG CCG AAA CCG CAG CCG CAG CCG CAG CCG CAG CCG AAA CCG CAG CCG AAA CCG GAA CCG GAA G; SEQ ID NO 12) and PX-low (5'TC GAC TTC CGG TTC CGG TTT CGG CTG CGG TTT CGG CTG CGG CTG CGG CTG CGG CTG CGG TTT CGG CTG CGG CTG CGG C; SEQ ID NO 13) oligonucleotides encoding the plus and minus strands of the duplex and contain XhoI and Sall cohesive ends on 5' and 3' end, respectively. The PX duplex is ligated into XhoI linearized, dephosphorylated pSC6H pDC6H and pFC6H plasmids, to generate the expression vectors pSPXC6H (DSM 11236), pDPXC6H and pFPXC6H, respectively. The final constructs are verified by dideoxynucleotide sequencing using Sequenase 2.0 (United States Biochemical).

Example 3: Expression and purification of pentabodies.

Pab fusion proteins are expressed in *E. coli* SG13009. The cultures are grown on a shaker at 37°C up to OD₆₀₀ ~0.5, then 1 mM IPTG is added to induce protein synthesis, followed by

further 4 h incubation at 30°C. Bacteria are collected by centrifugation (8000 x g, 15 min, 4°C) and frozen at -70°C. Bacterial pellet is resuspended in PBS pH 7.4, 1mM EDTA, 1mg/ml lysozyme, incubated for 30min at room temperature and subjected to three rounds of freezing/thawing (liquid nitrogen/37°C). The lysate is incubated for 15 min at RT with 0.1mg/ml of DNase I and after centrifugation (23,000 x g, 15 min, 4°C) the supernatant is collected. Imidazol is added to a final concentration of 5 mM and the recombinant protein is absorbed on 2ml of Ni-NTA resin (QIAGEN), equilibrated in 5 mM imidazol, PBS pH 7.4. After extensive wash with PBS containing 5 mM and 20 mM imidazol, retained proteins are eluted with PBS containing 250 mM imidazol. After extensive dialysis against PBS, 1 mM EDTA, proteins are concentrated 5 times with Centriprep 10 concentrator (Amicon), aliquoted and stored at -20°C.

The Pab-S and Pab-F fusion proteins are expressed at high levels (> 30 mg/l), allowing efficient one step metal-chelating affinity chromatography under native conditions. The amount of soluble Pab-D protein is lower than that of Pab-S and Pab-F.

Affinity purified Pab-S and Pab-F molecules are fractionated by FPLC gel filtration. A single elution peak corresponding to a protein of about 85 kDa is observed for Pab-S, whereas two major elution peaks corresponding to the proteins of about 90 and 180 kDa are observed for Pab-F. The high molecular weight peak presumably resulted from dimer formation *via* unsaturated cystein present in peptide F. In both cases, fractions of the elution peak corresponding to a 85-90 kDa protein are collected, pooled and used for binding studies.

Example 4: Equilibrium binding studies.

Cell dilution experiments and homogenous competition curves (i.e. competition for the cell binding between the labeled and the unlabeled forms of the same ligand) are described by a monovalent equilibrium model. The free (F) and bound (B) radioactivity is corrected from the non-immunoreactive fraction (NIF) and non-specific binding (NSB) respectively. Both are expressed in mol/l, taking into account the specific activity of each isotopic dilution (B and F, respectively). The parameters Kd, R, NSB and NIF, where R is the molarity of binding sites and Kd is the dissociation equilibrium constant, are fitted by non-linear regression of the corrected Scatchard equation

$$\frac{B - NSB}{F - NIF} = \frac{R}{Kd} - \frac{(B - NSB)}{Kd}$$

to the experimental data.

To determine maximal immunoreactivity, ^{125}I -labeled Pab-S, B1 IgG or B1 Fab' (20 nCi) are incubated with serial dilutions of freshly harvested BCL1 cells (0.3 to 100×10^5 cells per 100 μl). For competition assays, the labeled compounds are incubated with serial dilution of unlabeled competitors and BCL1 cells (25, 50 or 100×10^5 per 100 μl). Comparative experiments are performed the same day with the same batch of cells, in 96-wells plates, in triplicates, in a final volume of 150 μl of PBS supplemented with 1mg/ml of Bovine Serum Albumin (BSA), at 4°C, under agitation and for 2.5 h. After centrifugation, the amounts of both free and bound ^{125}I are measured as above from an aliquot of the supernatant and of the pellet washed once with PBS (4°C).

In preliminary solid phase competition assay Pab-S is found to compete the binding of ^{125}I labeled B1 IgG to BCL1 IgM idotype coated on plastic. However, since spatial arrangement of target molecule on the surface may influence multivalent binding, the equilibrium binding parameters of Pab-S are determined directly on live BCL1 cells, which represent a more relevant biological surface. Purified Pab-S is labeled with ^{125}I and is shown to bind to BCL1 idotype on the cell surface. It can be competed by unlabeled Pab-S, B1 IgG, and by a much higher concentration of peptide S, but not by the control Pab-F pentamer. The competitions of ^{125}I -labeled Pab-S binding by unlabeled Pab-S at 3 different cell concentration and a cell dilution experiment are used together for curve fitting to obtain the equilibrium binding parameters.

The apparent equilibrium binding constant of Pab-S is found to be similar to that of B1 IgG (around 1nM) which represents a 2×10^5 fold increase in avidity compared to the peptide S itself (IC_{50} around 200 μM). As control, the same binding experiments are performed with ^{125}I labeled B1 IgG and B1 Fab' fragment. The obtained constants are consistent with already published data. The Scatchard equilibrium binding constants are:

Fragment	Kd [nM]
Pab-S	1.0 ± 0.4
IgG	0.8 ± 0.4

Fab'	3.5 ± 0.1
------	---------------

The lower molarity of binding sites found for Pab-S compared to B1 IgG and Fab provides evidence for a multivalent nature of Pab-S binding.

In another series of experiments we compare the capacity of soluble BCL₁ IgM to compete the binding of either Pab-S or B1 IgG to BCL₁ cells. The results showed that much higher concentration of soluble BCL₁ IgM is needed to compete for Pab-S as compared to B1 IgG (100 vs 4 nM for 80% displacement). These data indicate a higher avidity of Pab-S for the cell surface-immobilized BCL₁ IgM, compared to the soluble BCL₁ IgM. Importantly, no inhibition of Pab-S binding to BCL₁ IgM idiotype is observed in the presence of up to 30 % (v/v) of human or mouse BALB/c serum.

Example 5: Biochemical characterization of Pab-S chimeric protein.

The concentration of purified Pab-S is determined by the method of Waddell, based on an absorption difference at 215 and 225 nm as well as by Bradford protein assay (Bio-Rad). Pab-S molecule is characterized using FPLC gel filtration, SDS-PAGE and CD spectrometry. Oxidized form of Pab-S with inter-chain disulfide bonds is formed by air oxidation during the purification procedure and dialysis against PBS. Completely reduced form is obtained by incubation with 100 mM DTT at 37°C for 30 min, followed by extensive dialysis against PBS, 1mM EDTA, 1mM β-mercaptoethanol.

Under non-denaturing conditions, a single elution peak, corresponding to a protein of about 85 kDa is detected for both oxidized and reduced forms of Pab-S after gel filtration on Superdex G200 column (equilibrated in PBS, 1 mM EDTA, ± 1mM β-mercaptoethanol, elution is monitored at 280 nm with an integrated UV detector (the proteins are analyzed on 10-15% gradient). Under denaturing conditions (Pab-S is denatured by boiling for 20 min in 4 M urea and 100 mM DTT and renatured by extensive dialysis against PBS, 1mM EDTA, 1mM β-mercaptoethanol) in SDS-PAGE, a major protein with apparent molecular weight of about 85 kDa is observed for the oxidized form of Pab-S molecule, whereas a protein with apparent molecular weight of about 17 kDa is observed for the reduced form of Pab-S, in agreement with a covalent pentameric structure. A minor band with apparent mobility of about 68 kDa observed for the oxidized Pab-S corresponds to a partially reduced pentamer,

where only four chain out of five are covalently link by disulfide bridges. From the analysis of the CD spectrum of Pab-S pentamer an α -helical content of about 54 % is determined by fitting the experimental data with a set of reference proteins as described in Vogel (Biochemistry (1987), 26, 4562-72). This is in a good agreement with the length of the α -helical coiled-coil domain within the entire molecule. Taken together these data indicate that Pab-S molecule is a stable homopentamer of about 85 kDa with a monomer subunits of about 17 kDa held together by in α -helical coiled-coil bundle, where the five chains are covalently crosslinked by disulfide bonds.

It is known that coiled coil structures can undergo reversible denaturation. Indeed, the Pab-S molecule is denatured in urea at 95°C under reducing conditions and refolded into the pentamer by simple dialysis against PBS, as demonstrated by FPLC gel filtration of renatured Pab-S labeled with ^{125}I . Noteworthy, renaturation of Pab-S restores full binding activity as shown by binding competition on BCL1 cells.

To visualize the spatial arrangement of Pab-S molecular modeling of its three-dimensional structure is undertaken. Pab-S molecule is modeled using the computer graphics program TOURBO-FRODO (Roussel & Cambillan (1989) in *Silicon Graphics Geometry Partner Directory (Fall 1989)*, eds. Silicon Graphics (Silicon Graphics, Mountain View, CA), pp.77-78). The structure is further refined by the X-PLOR program version 3.1 (Brunger. (1992) in X-PLOR version 3.1 A system for X-ray Crystallography and NMR, (New Haven, Yale University Press)) using the following procedure: a 5 ps molecular dynamics simulation at 300 K and then 1,000 steps of conjugate gradient minimization, carried out with 1/r dependent dielectric constant.

The stereochemical analysis shows that frequently occurring type I conformation of β -turn allows formation of disulfide bond between the two cysteins of peptide S. This conformation is chosen, as the most likely, for the peptide ligand. For the hinge region a molecular dynamic simulation with subsequent energy minimization resulted in a set of possible conformations. Out of it, a "good-looking" conformation is arbitrary chosen. Finally, previously modeled five-stranded α -helical coiled-coil structure is taken for the pentamerization domain. The analysis suggests a fivefold symmetrical structure of Pab-S protein. The analysis of the modeled Pentabody structure shows that such a molecule

should be capable of simultaneous binding to five surface receptor, provided they are located up to 80 Å apart from each other. This criterion is largely satisfied in the case of the surface immunoglobulin receptor, whose variable domains can be spaced at about 50 Å, judging by their van der Waals contact radii.

Example 6: Use of pentabody in *in vitro* for assays

To measure the level of an antigen of interest such as prostate specific antigen (PSA) which is an enzyme of the kallikrein family, peptides displayed on a phage display library may be screened for those binding specifically the antigen of interest, such as PSA, with a great specificity and avidity. Screening may be as described in the examples above, particularly example 1. Pentabody expressing the selected peptide(s) in pentameric form may be constructed, expressed and purified as described in the examples above, particularly examples 2 and 3. The selected pentabodies may be labeled with a marker enzyme such as peroxidase.

Antibodies specific for the antigen of interest, such as PSA, may be immobilized on a solid phase such as plates or polystyrene balls. The immobilized, unsolubilized antibodies may be incubated with the test fluid, such as serum or other biological fluid for the first step of immunoabsorption. After incubation, the solid phase may be washed.

The selected pentabody labeled with a marker enzyme may be incubated with the immunoabsorbed antigen, such as PSA. Then the substrate for the marker enzyme conjugated to the pentabody and a chromogen may be added. The optical density may then be measured. The optical density of the chromogen will be proportional to the amount of antigen, such as PSA, present. The assays of the invention may also comprise radioimmunoassays using the pentabodies of the invention, the above approach and methods known in the art.

Deposition of Microorganisms

The following microorganisms are deposited according to the Budapest Treaty with the DSM - Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig:

E. coli SG13009 pSXC6H

21.10.1996

DSM11236

Sequenzlisting

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: CIBA-GEIGY AG
- (B) STREET: Klybeckstr. 141
- (C) CITY: Basel
- (E) COUNTRY: Switzerland
- (F) POSTAL CODE (ZIP): 4002
- (G) TELEPHONE: +41 61 696 11 11
- (H) TELEFAX: + 41 61 696 79 76
- (I) TELEX: 962 991

(i) APPLICANT:

- (A) NAME: University of Lausanne
- (B) STREET: Chemin des Boveresses 155
- (C) CITY: Epalinges
- (E) COUNTRY: Switzerland
- (F) POSTAL CODE (ZIP): 1066
- (G) TELEPHONE:
- (H) TELEFAX:
- (I) TELEX:

(i) APPLICANT:

- (A) NAME: Swiss Institute for Allergy and Astma Research
- (B) STREET: Obere Stasse 22
- (C) CITY: Davos
- (E) COUNTRY: Switzerland
- (F) POSTAL CODE (ZIP): 7270
- (G) TELEPHONE:
- (H) TELEFAX:
- (I) TELEX:

(i) APPLICANT:

- (A) NAME: ISREC
- (B) STREET: Chemin des Boveresses 155
- (C) CITY: Epalinges

- (E) COUNTRY: Switzerland
- (F) POSTAL CODE (ZIP): 1066
- (G) TELEPHONE:
- (H) TELEFAX:
- (I) TELEX:

(ii) TITLE OF INVENTION: Oligomers

(iii) NUMBER OF SEQUENCES: 13

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ser	Val	Trp	Arg	Trp	Leu	Pro	Tyr	Asp	Lys	Tyr	Glu
1				5					10		

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 21 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ala Asp Gly Ala Cys Arg Asn Pro Trp Cys
1 5 10

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Thr Ala Ala Gly Leu Cys Glu Phe Asp Gln Ala Leu Leu Arg Tyr Thr
1 5 10 15

Cys Pro Thr

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

AGATCCTCGA GGGTGGAGAC TGCTGCCCAC AGATGCTTAG A

41

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- 22 -

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GCACTAGTAG AACCGCCACC CGGGGT

26

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GATCCGCTGA CGGCGCTTGC CGTACCCCGT GGTGCGGTC

39

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TCGAGACCGC ACCACGGGTT ACGGCAAGCG CCGTCAGCG

39

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GATCCACTGC TGCAGGTCTG TCGAATCCG ACCAGC

36

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TCGAGCTGGT CGAATTCGCA CAGACCTGCA GCAGTG

36

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GATCCTCTGT TTGGCGTTGG CTGCCGTACG ACAAATACGA AC

42

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

TCGAGTTCGT ATTTGTCGTA CGGCAGCCAA CGCCAAACAG AG

42

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

TCGAGCGGCA GCCGCAGCCG AAACCGCAGC CGCAGCCGCA GCCGCAGCCG AAACCGCAGC

60

CGAAACCGGA ACCGGAAG

78

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

TCGACTTCCG GTTCCGGTTT CGGCTGCGGT TTCGGCTGCG GCTGCGGCTG CGGCTGCGGT 60

TTCGGCTGCG GCTGCGGC 78

Claims

1. An oligomer comprising 2 or more than 2 units, wherein each unit comprises a peptidic domain capable of oligomerizing and a domain capable of binding to an acceptor, wherein the oligomerizing domain is not an antibody or a functional antibody fragment from the constant region.
2. An oligomer according to claim 1 comprising more than 4 units.
3. An oligomer according to claim 1 consisting of 5 units.
4. An oligomer according to claim 1, wherein the acceptor is a antibody or a receptor.
5. An oligomer according to claim 1, wherein each of said units has less that 600 amino acids.
6. An oligomer according to claim 1, wherein the peptidic domain capable of oligomerizing and the domain capable of binding to an acceptor are connected via a spacer (hinge region).
7. An oligomer according to claim 6, wherein the spacer comprises a proline-rich region.
8. An oligomer according to claim 1, wherein at the C-terminus of some or all units a further functional domain is attached.
9. An oligomer according to claim 1, wherein the individual units oligomerize spontaneously.
10. Use of the pentamerization domain of the Cartilage Oligomeric Matrix Protein for the pentamerization of low molecular weight compounds or peptides that are not part of the Cartilage Oligomeric Matrix Protein.
11. An oligomer according to claim 1, wherein the oligomerizing domain is the pentamerization domain of the Cartilage Oligomeric Matrix Protein.
12. An oligomer according to claim 1 comprising units that bind to distinct acceptors.
13. A unit capable of oligomerizing according to claim 1.

14. Use of an oligomer according to claim 1 for identification and/or marking of acceptors.
15. Use of an oligomer according to claim 1 for eukaryotic cell, bacteria or viruses targeting.
16. Use of an oligomer according to claim 1 for cell targeting.
17. Use according to claim 16 for targeting of B-cell lymphomas.
18. Use of an oligomer according to claim 1 for the inhibition of protein-protein interactions.
19. Use of an oligomer according to claim 12 as a chelating agent.
20. Use of an oligomer according to claim 12 as a crosslinking agent.
21. Use of oligomers according to claim 1 for the construction of libraries.
22. Use of oligomers according to claim 1 for the induction of apoptosis.
23. Use of oligomers according to claim 1 for the intracellular inhibition of transcription factor binding, gene regulating molecules and/or enzymatic activities.
24. Use of oligomers according to claim 1 for prevention of tumor metastatization.
25. Use of oligomers according to claim 1 *in vitro* as one of the binding reagents in an enzyme immunoassay.
26. Use of oligomers according to claim 1 *in vitro* as one of the binding reagents in radioimmunoassays.
27. Method for the synthesis of a unit according to claim 13.
28. An expression vector for the synthesis of a unit according to claim 13.
29. A host, comprising an expression vector for the synthesis of a unit according to claim 13.
30. A microbiological host, comprising an expression vector for the synthesis of a unit according to claim 13.
31. A method for the production of an oligomer according to claim 1.

32. An oligomer according to claim 1, wherein the epitopes are used for vaccine development.

DECLARATION AND POWER OF ATTORNEY FOR UNITED STATES PATENT APPLICATION

☒ Original

☐ Supplemental

☐ Substitute

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a United States patent is sought on the invention entitled

METHOD FOR THE OLIGOMERISATION OF PEPTIDES

the specification of which:

☐ is attached hereto.

☐ was filed on _____ as Application No. _____
(day/month/year)

and, if this box (☐) contains an *

☐ was amended on _____
(day/month/year)

☒ was filed as Patent Cooperation Treaty international Application No.

PCT/EP 97/05897 on October 24, 1997
(day/month/year)

and, if this box (☐) contains an *

☐ entered the national stage in the United States and was accorded Application No. _____

and, if this box (☐) contains an *

☐ was amended, subsequent to entry into the national stage, on _____
(day/month/year)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above and, if this application was filed as a Patent Cooperation Treaty international application, by any amendments made during the international stage (including any made under Patent Cooperation Treaty Rule 91, Article 19 and Article 34).

I acknowledge my duty to disclose all information which is known by me to be material to the patentability of this application as defined in 37 C.F.R. § 1.56.

I hereby appoint the registered practitioners associated with Customer No. 001095, respectively and individually, as my attorneys and agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

If these brackets contain an X [X], I hereby authorize the registered practitioners associated with Customer No. 001095 and any others acting on my behalf to take any action relating to this application based on communications from the Patents and Trademarks Division of Novartis Services AG, Basle, Switzerland, or an affiliate thereof or a successor thereto, without direct communication from me.


Please address all communications to Michael W. Glynn, Novartis Corporation, Patent and Trademark Department, 564 Morris Avenue, Summit, NJ 07901-1027.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full name of sole
or first joint inventor

Alexey TERSKIKH

Inventor's signature



Date

May 17 99
(day/month/year)

Residence

Lausanne, Switzerland CHX

Citizenship

Russia

Post Office Address


Chemin des Begonias
1018 Lausanne
Switzerland

IMPORTANT: Before this declaration is signed, the patent application (the specification, the claims and this declaration) must be read and understood by each person signing it, and no changes may be made in the application after this declaration has been signed.

Full name of second
joint inventor, if any

Reto CRAMERI

Inventor's signature



Date

01.05.1999

(day/month/year)

Residence

Davos Platz, Switzerland

CHX

Citizenship

Switzerland

Post Office Address

**Edenstrasse 13
7270 Davos Platz
Switzerland**

Full name of third
joint inventor, if any

Jean-Pierre MACH

Inventor's signature

Date

(day/month/year)

Residence

Lausanne, Switzerland

Citizenship

Switzerland

Post Office Address

**Av. de Valmont 20
1010 Lausanne
Switzerland**

Full name of fourth
joint inventor, if any

Jean-Marc LE DOUSSAL

Inventor's signature

Date

(day/month/year)

Residence

Paris Cedex, France

Citizenship

France

Post Office Address

20 Rue Constantinople

**75008 Paris Cedex
France**



Full name of second
joint inventor, if any

Reto CRAMERI

Inventor's signature

Date

(day/month/year)

Residence

Davos Platz, Switzerland

Citizenship

Switzerland

Post Office Address

**Edenstrasse 13
7270 Davos Platz
Switzerland**

Full name of third
joint inventor, if any

Jean-Pierre MACH

Inventor's signature

Jean-Pierre Mach

Date

27 April 1999
(day/month/year)

Residence

Lausanne, Switzerland

Citizenship

Switzerland

Post Office Address

**Av. de Valmont 20
1010 Lausanne
Switzerland**

Full name of fourth
joint inventor, if any

Jean-Marc LE DOUSSAL

Inventor's signature

Date

(day/month/year)

Residence

Paris Cedex, France

Citizenship

France

Post Office Address

20 Rue Constantinople

**75008 Paris Cedex
France**

Full name of second
joint inventor, if any

Reto CRAMERI

Inventor's signature

Date

(day/month/year)

Residence

Davos Platz, Switzerland

Citizenship

Switzerland

Post Office Address

**Edenstrasse 13
7270 Davos Platz
Switzerland**

Full name of third
joint inventor, if any

Jean-Pierre MACH

Inventor's signature

Date

(day/month/year)

Residence

Lausanne, Switzerland

Citizenship

Switzerland

Post Office Address

**Av. de Valmont 20
1010 Lausanne
Switzerland**

Full name of fourth
joint inventor, if any

Jean-Marc LE DOUSSAL

Inventor's signature

Date

23 April 1999
(day/month/year)

Residence

Paris Cedex, France

FR

Citizenship

France

Post Office Address

**20 Rue Constantinople
75008 Paris Cedex
France**

Full name of fifth
joint inventor, if any

⁵ Andrey KAJAVA

Inventor's signature

Kajava

Date

11/06/99

(day/month/year)

Residence

Lausanne, Switzerland

CAK

Citizenship

Russia

Post Office Address

20, chemin du Mont Tendre
1007 Lausanne
Switzerland

11/06/99 11:06:00